

*AHELP for CIAO 3.4*

pix_get_energy

Context: [pixlib](#)*Jump to:* [Description](#) [Bugs](#) [See Also](#)

Synopsis

Return the photon energy corresponding to a given grating dispersion (GAC) value.

Syntax

```
Double_Type pix_get_energy( Double_Type x, Double_Type y )
```

Description

Calculates the photon energy, in keV, for a given grating position and using the current `pixlib` settings. The `x` and `y` values are in the GAC coordinate system, and so represent the grating dispersion and cross-dispersion angles in degrees. The grating order – as set by `pix_set_grating()` – must be non-zero for this routine to work.

The `pix_get_grating_wavelength()` routine can be used to find the wavelength of the photon instead of the energy.

Bugs

See the [bugs page for the pixlib library](#) on the CIAO website for an up-to-date listing of known bugs.

See Also

modules[pixlib](#)*pixlib*[pix_gac to gdp](#), [pix_gdp to gac](#), [pix_get_grating_wavelength](#)

The Chandra X-Ray Center (CXC) is operated for NASA by the Smithsonian Astrophysical Observatory.
60 Garden Street, Cambridge, MA 02138 USA.
Smithsonian Institution, Copyright © 1998–2006. All rights reserved.

URL:
http://cxc.harvard.edu/ciao3.4/pix_get_energy.html

Last modified: December 2006

